

Lens spaces given from L-space homology 3-spheres

Motoo Tange

Department of Mathematics, Osaka University,
Machikaneyama 1-1 Toyonaka, Osaka 560-0043, Japan
e-mail: tange@math.sci.osaka-u.ac.jp

Abstract

We consider the problem when lens spaces are given from homology spheres, and demonstrate that many lens spaces are obtained from L-space homology sphere which the correction term $d(Y)$ is equal to 2. We show an inequality of slope and genus when Y is L-space and $Y_p(K)$ is lens space. ¹ ²

1 Introduction

In the present paper we define lens space $L(p, q)$ to be the p/q -Dehn surgery of unknot, where p, q are coprime integers. Note that this orientation is opposite to usual one. Let $K \subset S^3$ be a knot whose Dehn surgery is homeomorphic to a lens space. Then we call that K admits *lens surgery on S^3* or simply *lens surgery*. The main problems of lens surgery involve when a lens space is obtained from Dehn surgery of a classical knot or when a knot K admits lens surgery.

The Alexander polynomial $\Delta_K(t)$ of classical knot K is useful for lens surgery problem. For example for a certain knot K yielding a lens space if $\Delta_K(t)$ is trivial, then K is unknot, since the degree of Alexander polynomial coincides with the genus of K . If $\Delta_K(t)$ is $t - 1 + t^{-1}$, then K is the trefoil knot in [3]. Although it is unknown whether the uniqueness holds otherwise, it seems that Alexander polynomial is an effective invariant for lens surgery. As showed in [17], the doubly primitive knots which yields $L(p, q)$ can be distinguished by the Alexander polynomials.

Many people have been done research on condition for lens surgery. In [1] J. Berge has defined doubly primitive knots, each of which is a classical knot yielding lens space. Conversely he conjectured that these knots are all knots admitting lens surgery. He had fallen the doubly primitive knots into several types. But unfortunately the classification is not sure whether it is complete. If the uniqueness above holds, then his conjecture is solved.

P. Ozsváth and Z. Szabó have proven the constraints of Alexander polynomial $\Delta_K(t)$ of knots yielding lens space by using Heegaard Floer homology. The formula in Section 10.3 or Theorem 7.2 in [8] represents the relationship between correction term and Alexander polynomial. This formula is due to the surgery exact triangle among S^3 , $S_0^3(K)$ and $L(p, q)$.

In [2] R. Fintushel and R. Stern have shown a criterion for obtaining a lens space from a Dehn surgery of a homology sphere. The statement is the following.

¹Keyword: lens surgery, Heegaard Floer homology, Alexander polynomial, Homology sphere

²MSC: 57M25, 57M27, 57R58

Theorem 1.1 ([2]) *The lens space $L(p, q)$ can be obtained as integral positive surgery on a knot in an integral homology three-sphere if and only if q is a square modulo p .*

On the other hand P. Ozsváth and Z. Szabó have shown that for positive integers k not divisible by 4, $L(2k(3 + 8k), 2k + 1)$ cannot be obtained from Dehn surgery of S^3 , but can be obtained from Dehn surgery of a Brieskorn homology sphere and moreover the lens space is obtained from 2-component link. The detailed proof is in [8].

We would like to consider when a lens space can be obtained from a Dehn surgery of a knot in a homology sphere. This issue seems to be worth considering but it does not seem that such studies about Dehn surgery of homology spheres have been developed remarkably. This present paper attends to Dehn surgeries of L-space homology spheres, which is defined as the manifolds whose Heegaard Floer homology is isomorphic to that of S^3 . The Heegaard Floer theoretic arguments of Dehn surgery of L-space homology sphere are the same as S^3 up to the absolute grading. They lead to the three results: the surgery formula between the correction term and the Alexander polynomial (Equation (4)), the constraint of Alexander polynomial of lens surgery knot (Lemma 3.1), and the lower bound of the slope by the genus of the knot. The lower bound is

$$2g(K) - 1 \leq p.$$

Here we state an upper bounds, where it generalizes the main theorem in [12]. The notation $d(Y)$ is the correction term of Y , whose definition is in [8].

Theorem 1.2 *Let Y be an L-space homology sphere. Suppose that $Y_p(K)$ is a lens space and K is a non-trivial knot in Y . Then $g(K) + 2d(Y) > 0$ and the following bound holds:*

$$p < \frac{4g(K)(g(K) + 1)}{g(K) + 2d(Y)}. \quad (1)$$

We will prove the proof in Section 2.

By using the lower and upper bounds, we can understand that as $d(Y)$ is more increasing, $g(K)$ must also become more increasing. Note that lens spaces coming from L-space homology spheres are not empty other than S^3 , for $L(22, 3)$ is the 22-Dehn surgery of Poincaré homology sphere, which the correction term is 2. When $Y = S^3$, this theorem means Theorem 1 in [12].

In the case where an L-space homology sphere Y is $d(Y) = 2$, Y can give many lens spaces as demonstrated later on. But when let $d(Y) \neq 0, 2$, there do not exist such lens spaces in the range of $p < 1000$ and $|d(Y)| \leq 40$ by Maple's computation. For example Y is $\Sigma(2, 3, 5) \# \Sigma(2, 3, 5)$. It is not likely that lens space can be constructed from reducible manifolds. We conjecture the following here.

Conjecture 1.1 *Let Y be any irreducible L-space homology sphere with $d(Y) \neq 0, 2$. Then there never exist knot $K \subset Y$ such that $Y_p(K)$ is a lens space for a positive integer p .*

The author proved this conjecture in the case where Y is Poincaré homology sphere with reverse orientation, namely $d(Y) = -2$, (see [15]).

In Section 4 we illustrate many lens spaces are obtained from L-space homology sphere with $d(Y) = 2$. In practice some of lens spaces are constructed from Poincaré homology sphere $\Sigma(2, 3, 5)$ along knots whose dual knots are 1-bridge knots in the lens spaces. We put a table of such lens spaces up to $p \leq 2007$ in the last of the Section. The L-space homology sphere with $d(Y) = 2$ gives several infinite sequences of lens spaces except fewer examples. In particular from Poincaré homology sphere infinite many lens spaces can be constructed by Dehn surgeries.

The author does not know whether there exist lens spaces which are given rise to from both S^3 and $\Sigma(2, 3, 5)$. Assume that exists such a lens space $L(p, q)$, the two

knots K, K' in $S^3, \Sigma(2, 3, 5)$ respectively. Then p is odd and the genus of K' is $\frac{p+1}{2}$. Moreover the Alexander polynomial of the two knots must have the following relation $\Delta_{K'}(t) = \Delta_K(t) - (t^{\frac{p-1}{2}} + t^{-\frac{p-1}{2}}) + (t^{\frac{p+1}{2}} + t^{-\frac{p+1}{2}})$, by using Equations (4) as explained later on.

2 The exact triangle and the Alexander polynomial

From this section on, we denote Y as an L-space homology sphere. If a positive p -Dehn surgery $Y_p(K)$ along knot K is a lens space, then we have the short exact sequences for any $0 \neq i \in \mathbb{Z}/p\mathbb{Z}$,

$$0 \rightarrow \bigoplus_{j \equiv i \pmod p} HF^+(Y_0, j) \rightarrow HF^+(Y_p(K), Q(i)) \rightarrow HF^+(Y) \rightarrow 0 \quad (2)$$

and for $i = 0$,

$$0 \rightarrow HF^+(Y) \rightarrow \bigoplus_{j \equiv 0 \pmod p} HF^+(Y_0, j) \rightarrow HF^+(Y_p(K), Q(i)) \rightarrow 0. \quad (3)$$

From this exact sequences for any i the following formula is induced as in [8]:

$$d(Y) - d(Y_p(K), Q(i)) + d(L(p, 1), i) = 2t_i(K), \quad (4)$$

where $Q(i) = hi + c$, h is the homology class of the dual knot K^* of K and $c = \frac{(h+1+p)(h-1)}{2}$ by the result in [16]. The each integer $t_i(K)$ is the i -th Turaev torsion of $Y_0(K)$. It is positive when $Y_p(K)$ is L-space.

Here we use the identifications $\text{Spin}^c(L(p, q)) \cong H_1(L(p, q)) \cong H^2(L(p, q)) \cong \mathbb{Z}/p\mathbb{Z}$. By taking the summation of Equation (4) over $i \in \mathbb{Z}/p\mathbb{Z}$, we obtain

$$\begin{aligned} p(d(Y) + 2\lambda(L(p, q)) - 2\lambda(L(p, 1))) &= 2 \sum_{i \in \mathbb{Z}} t_i(K) \\ &= 2 \sum_{i \in \mathbb{Z}} i^2 a_i(K) \\ &= \Delta_K''(t)|_{t=1}, \end{aligned} \quad (5)$$

where the Alexander polynomial is normalized and symmetrized and Casson-Walker invariant λ is computed by the Rustamov's formula in [13]:

$$\sum_{\mathfrak{s} \in \text{Spin}^c(W)} \left(\chi(HF_{red}(W, \mathfrak{s})) - \frac{1}{2}d(W, \mathfrak{s}) \right) = |H_1(W, \mathbb{Z})| \lambda(W),$$

where W is any rational homology sphere.

By the same argument as the proof of Theorem 7.2 in [8] the statement replaced S^3 with Y in that theorem also holds.

We can compute the coefficients of Alexander polynomial of the knot yielding lens surgery as follows. Let $[[\alpha, \beta]]$ be $[\alpha, \beta] \cap \mathbb{Z}$ and $[\gamma]_p$ be a reduction modulo p of γ with $0 \leq [\gamma]_p < p$. We define $\Phi_{p,q}^k(h)$ as $\#\{j \in [[1, h']][qj - k]_p \in [[1, h]]\}$, where $h = [h]_p, h' = [h^{-1}]_p$. For any class $i \in \mathbb{Z}/p\mathbb{Z}$ the reduced coefficient $\tilde{a}_i(K)$ of the Alexander polynomial of knot K is defined as $\sum_{j \equiv i \pmod p} a_j(K)$. When $L(p, q) = Y_p(K)$, $\tilde{a}_j(K)$ is computed as follows:

$$\tilde{a}_i(K) = -m + \Phi_{p,q}^{hi+c}(h). \quad (6)$$

3 The upper bound of the slope by the Seifert genus

In this section we prove Theorem 1.2. First, we introduce the two lemmas below.

Lemma 3.1 *Let Y be an L -space homology sphere. If $Y_p(K)$ is a lens space, the degree of the Alexander polynomial $\Delta_K(t)$ coincides with the Seifert genus $g(K)$.*

Lemma 3.2 *Let Y be L -space homology sphere and $Y_p(K)$ be L -space. Then $\Delta_K(t)$ has the form*

$$\Delta_K(t) = (-1)^k + \sum_{j=1}^k (-1)^{k-j} (t^{n_j} + t^{-n_j})$$

for some increasing sequence of positive integers $0 < n_1 < n_2 < \dots < n_k$.

The proofs of these lemmas' are immediately derived from an application to [9] and [10]. Note that the property of Lemma 3.2 is also preserved for the reduced polynomial $\sum_{|i| < \frac{p}{2}} \tilde{a}_i(K) t^i$ when $g(K) < \frac{p}{2}$ or $\sum_{|i| < \frac{p}{2}} \tilde{a}_i(K) t^i + t^{\frac{p}{2}} + t^{-\frac{p}{2}}$ when $g(K) = \frac{p}{2}$.

From this assumption $Y - K$ is irreducible and from the main theorem in [18] and Lemma 3.2, K is a fiber knot. This leads to the result of Lemma 3.1.

For the proof of Theorem 1.2 we use the following proposition in [12]. Note that λ is multiplied by $-1/2$ for the Casson-Walker invariant in [12] and the definition of orientations of lens spaces is the opposite to the one in [12].

Proposition 3.1 (Proposition 2.4.[12]) *Suppose that L is a lens space with $|H_1(L)| = p$, and that*

$$2\lambda(L) - 2\lambda(L(p, 1)) \leq \frac{1}{4} \left(\frac{p}{4} - 1 \right).$$

Then L is homeomorphic to one of $L(p, 1)$, $L(p, 2)$, or $L(p, 3)$.

Proof of Theorem 1.2) From the Frøyshov's inequality in [12] and Equation (5),

$$\begin{aligned} 2p(\lambda(L(p, q)) - \lambda(L(p, 1))) &= 2 \sum_{i \geq 1} i^2 a_i(K) - pd(Y) \\ &\leq g(K)(g(K) + 1) - pd(Y). \end{aligned}$$

We assume that

$$p \geq \frac{4g(K)(g(K) + 1)}{g(K) + 2d(Y)}. \quad (7)$$

Then we have $g(K)(g(K) + 1) - pd(Y) \leq \frac{p}{4}(\frac{p}{4} - 1)$ by solving the quadratic inequality. By Proposition 3.1 $L(p, q) = L(p, 1)$, $L(p, 2)$, or $L(p, 3)$ hold. L -space homology sphere Y yielding such lens spaces must be $d(Y) = 0$ or 2 , due to Lemma 3.2 and the positivity of $t_i(K)$ for any i .

We consider the case of $Y_p(K) = L(p, 1)$. By Lemma 3.2 $d(Y) = 0$ or 2 . From $q = 1$ we have $h = h'$ and we may assume that $h < \frac{p}{2}$ by replacing h with $p - h$. Hence Formula (6) says $\tilde{a}_i = -m$ for some integer i . By Lemma 3.2 non-negative integer m is 0 or 1 . When $m = 0$, $h = h' = 1$ namely $\Delta_K(t) = 1 \bmod t^p - 1$ holds. K is trivial from Lemma 3.1 Therefore $Y = S^3$. When $m = 1$, $\tilde{a}_{-h'}(K) = -1 + h$ is $0, \pm 1$, or 2 by applying Lemma 3.2. Thus we have $h = 1, 2, 3$.

When $h = 1$, $h^2 = 1$ hence $p = 0$ and this is inconsistent.

When $h = 2$, $h^2 = 4 = p + 1$ hence $p = 3$. $\Delta_K(t) = 1 \bmod t^p - 1$.

When $h = 3$, $h^2 = 9 = p + 1$ hence $p = 8$, $\Delta_K(t) = t^{-4} - t^{-3} + t - 1 + t - t^3 + t^4$.

As a whole, the case of $L(p, q) = Y_p(K)$ classifies the following.

- $S_p^3(\text{trivial knot}) = L(p, 1)$ and $h = 1$.

- $Y_p(K_{1,p}) = L(p, 1)$, $d(Y) = 2$, p is odd, $h = 1$
and $\Delta_{K_{1,p}}(t) = t^{-\frac{p+1}{2}} - t^{-\frac{p-1}{2}} + 1 - t^{\frac{p-1}{2}} + t^{\frac{p+1}{2}}$.
- $Y_8(K_2) = L(8, 1)$, $d(Y) = 2$, $h = 3$, $\Delta_{K_2}(t) = t^{-4} - t^{-3} + t^{-1} - 1 + t - t^3 + t^4$.

In case of $Y_p(K) = L(p, 2)$ or $L(p, 3)$, analogous computation of $\tilde{a}_i$ and Lemma 3.1, 3.2 classifies the follows.

- $Y_7(K_3) = L(7, 2)$, $d(Y) = 0$ or 2
and $\Delta_{K_3}(t) = t^{-1} - 1 + t$ or $t^{-4} - t^{-3} + t^{-1} - 1 + t - t^3 + t^4$ respectively.
- $Y_{11}(K_4) = L(11, 3)$, $d(Y) = 0$ or 2 , $h = 5$ and $\Delta_{K_4}(t) = t^{-2} - t^{-1} + 1 - t + t^2$
or $t^{-6} - t^{-5} + t^{-2} - t^{-1} + 1 - t + t^2 - t^5 + t^{-6}$ respectively.
- $Y_{13}(K_5) = L(13, 3)$, $d(Y) = 0$ or 2 , $h = 5$ and $\Delta_{K_5}(t) = t^{-3} - t^{-2} + 1 - t^2 + t^3$
or $t^{-7} - t^{-6} + t^{-3} - t^{-2} + 1 - t^2 + t^3 - t^{-6} + t^{-7}$ respectively.
- $Y_{22}(K_6) = L(22, 3)$, $d(Y) = 2$, $h = 5$
and $\Delta_{K_6}(t) = t^{-11} - t^{-10} + t^{-6} - t^{-5} + t^{-2} - 1 + t^2 - t^5 + t^6 - t^{10} + t^{11}$.

Each of the cases does not satisfy Inequality (7). Therefore Inequality (1) is proven. $\square$

4 A table of several lens surgeries over Y with $d(Y) = 2$

In this section and the subsequent section Y is an L-space homology sphere with $d(Y) = 2$ for example Y is $\Sigma(2, 3, 5)$. Since lens surgery by K with $2g(K) - 1 = p$ includes the ones admitting lens surgery on S^3 we restrict our attention to lens surgery with $2g(K) - 1 < p$. We may be able to construct a lens space form both integral Dehn surgeries of S^3 and $\Sigma(2, 3, 5)$ but ignore such a lens space here. The homology class of the dual knot K^* of K is $h[l] \in H_1(L(p, q))$, where curve l is a core loop of a handlebody of genus one Heegaard splitting. The set of classes $\mathcal{H}(p, K) := \{\pm h^\pm\} \subset \mathbb{Z}/p\mathbb{Z}$ is an invariant of the lens surgery which is independent of choices of two handlebodies of the Heegaard decomposition and the orientation and we always consider any element in this set as the integers reduced to $\{1, 2, \dots, p\}$.

Any datum (p, q, h, g) in Table 1, and 2 represent lens space $L(p, q)$ and the minimal representative h in $\mathcal{H}(p, K)$ that the coefficient (6) satisfies Lemma 3.2, and $t_i(K)$ is all non-negative, and g is $\max\{i \in \{0, 1, \dots, \lfloor \frac{p}{2} \rfloor\} \mid -m + \Phi_{p,q}^{hi+c}(h) \neq 0\}$.

Let $L(p, q) = Y_p(K)$. We say that K is *0-bridge knot* if K is isotopic to a knot which lies on Heegaard surface of genus one Heegaard splitting, and *1-bridge knot* if the knot is non-0-bridge knot and is the union of two arcs embedded in the meridian disks of both handlebodies of genus one Heegaard splitting of the lens space. This definition is based on [1]. From any triplicity (p, q, h) we can uniquely find 0-bridge knot or 1-bridge knot having the datum.

Theorem 4.1 *Lens spaces $L(p, q)$ in Table 1 and 2 are constructed by p -Dehn surgery of knots in $\Sigma(2, 3, 5)$. Moreover the dual knots are 1-bridge knots in $L(p, q)$.*

Before the proof of this theorem we prove the following lemma.

Lemma 4.1 *Lens spaces in Table 1, 2 satisfy one of the list below for some $\ell \in \mathbb{Z} \setminus \{0\}$.*

- $p = 14\ell^2 + 7\ell + 1$, $h = \pm(7\ell + 2)^{\pm 1} \pmod{p}$, $2g = p + 1 - |\ell|$
- $p = 20\ell^2 + 15\ell + 3$, $h = \pm(5\ell + 2)^{\pm 1} \pmod{p}$, $2g = p + 1 - |\ell|$
- $p = 30\ell^2 + 9\ell + 1$, $h = \pm(6\ell + 1)^{\pm 1} \pmod{p}$, $2g = p + 1 - |\ell|$
- $p = 42\ell^2 + 23\ell + 3$, $h = \pm(7\ell + 2)^{\pm 1} \pmod{p}$, $2g = p + 1 - |\ell|$

- d') $p = 42\ell^2 + 47\ell + 13, h = \pm(7\ell + 4)^{\pm 1} \pmod p, 2g = p + 1 - |\ell|$
- e) $p = 52\ell^2 + 15\ell + 1, h = \pm(13\ell + 2)^{\pm 1} \pmod p, 2g = p + 1 - |\ell|$
- e') $p = 52\ell^2 + 63\ell + 19, h = \pm(13\ell + 8)^{\pm 1} \pmod p, 2g = p + 1 - |\ell|$
- f) $p = 54\ell^2 + 15\ell + 1, h = \pm(27\ell + 4)^{\pm 1} \pmod p, 2g = p + 1 - |\ell|$
- f') $p = 54\ell^2 + 39\ell + 7, h = \pm(27\ell + 10)^{\pm 1} \pmod p, 2g = p + 1 - |\ell|$
- g) $p = 69\ell^2 + 17\ell + 1, h = \pm(23\ell + 3)^{\pm 1} \pmod p, 2g = p + 1 - 2|\ell|$
- g') $p = 69\ell^2 + 29\ell + 3, h = \pm(23\ell + 5)^{\pm 1} \pmod p, 2g = p + 1 - 2|\ell|$
- h) $p = 85\ell^2 + 19\ell + 1, h = \pm(17\ell + 2)^{\pm 1} \pmod p, 2g = p + 1 - 2|\ell|$
- h') $p = 85\ell^2 + 49\ell + 7, h = \pm(17\ell + 5)^{\pm 1} \pmod p, 2g = p + 1 - 2|\ell|$
- i) $p = 99\ell^2 + 35\ell + 3, h = \pm(11\ell + 2)^{\pm 1} \pmod p, 2g = p + 1 - 2|\ell|$
- i') $p = 99\ell^2 + 53\ell + 7, h = \pm(11\ell + 3)^{\pm 1} \pmod p, 2g = p + 1 - 2|\ell|$
- j) $p = 120\ell^2 + 16\ell + 1, h = \pm(12\ell + 1)^{\pm 1} \pmod p, 2g = p + 1 - 2|\ell|$
- k) $p = 120\ell^2 + 20\ell + 1, h = \pm(20\ell + 2)^{\pm 1} \pmod p, 2g = p + 1 - 2|\ell|$
- l) $p = 120\ell^2 + 36\ell + 3, h = \pm(12\ell + 2)^{\pm 1} \pmod p, 2g = p + 1 - 2|\ell|$
- m) $p = 120\ell^2 + 104\ell + 22, h = \pm(12\ell + 5)^{\pm 1} \pmod p, 2g = p + 1 - |2\ell + 1|$
- n) $L(191, 34), h = 15$

Proof) It is only necessary to prove that data (p, q, h) given rise to by the 20 lists cover Table 1, 2 by direct computation. The computation of genus of K is due to Lemma 3.1.

□

Proof of Theorem 4.1) From each datum (p, q, h) in Lemma 4.1 we take the 1-bridge knot K^* in $L(p, q)$. The $-\tilde{a}_{-h'c-h'}(K)^*$ -surgery in the sense of T.Saito in [14] obtains a homology sphere Y . Thus there exists a knot $K \subset Y$ such that we have $Y_p(K) = L(p, q)$. The presentation of $\pi_1(Y)$ is

$$\left\langle x_1, x_2 \mid \prod_{i=1}^p x_1 x_2^{\delta_h(qi+1)}, \left(\prod_{i=1}^{h'-1} x_1 x_2^{\delta_h(qi+1)} \right) x_1 x_2^{-\tilde{a}_{h'c-h}} \right\rangle$$

by [17], where $\delta_h : \mathbb{Z}/p\mathbb{Z} \rightarrow \{0, 1\}$ is define to be $\delta_h(k) = 1$ when $k = 1, \dots, h \pmod p$, and $\delta_h(k) = 0$ otherwise.

By deforming this group presentation directly we can easily give an isomorphism $\pi_1(Y) \cong \langle x, y \mid (xy)^2 = x^3 = y^5 \rangle$. The right hand side is the same as the fundamental group of $\Sigma(2, 3, 5)$. Due to the celebrated resolution of Poincaré conjecture by G. Perelman in [11], Y is homeomorphic to $\Sigma(2, 3, 5)$.

□

The author would like to classify lens spaces obtained from Poincaré homology sphere. But the author could not control type n) in Lemma 4.1. If you go on calculating more, then a new sequence may be discovered.

We here illustrate two figures, which are Figure 1 and 2. Any point of Figure 1 represents a lens surgery over $\Sigma(2, 3, 5)$ with the slope $p \leq 2007$. The point (h, p) means that there exists a knot such that $L(p, q) = \Sigma(2, 3, 5)_p(K)$ and h is the minimal in the set $\mathcal{H}(p, K)$ defined in Section 4. Figure 2 represents hyperbolic lens surgeries over S^3 which are plotted in the order of slope from the smallest up to the same cardinality as plots in Figure 1. The horizontal and vertical axes mean the same as Figure 1. Compared with Figure 1 and 2 it follows that the existence of lens surgeries over $\Sigma(2, 3, 5)$ are localized near four quadratic functions. The right two of the them correspond to f), f'), g) and g'). To draw Figure 1, we referred to the last table of the article [1]. Here we give a rough conjecture on the basis of Figure 1.

Table 1: Lens spaces with $p \leq 711$ which homology sphere with $d(Y) = 2$ yield.

p	q	h	g	p	q	h	g	p	q	h	g
8	1	3	4	221	127	41	109	442	157	77	220
22	3	5	11	228	61	17	113	445	186	39	221
38	7	7	19	239	67	28	119	445	84	23	221
40	9	7	20	243	133	43	120	449	80	23	223
43	15	12	21	244	45	17	121	450	79	23	224
53	11	8	26	246	43	17	122	463	211	40	229
67	14	9	33	247	134	58	123	469	107	24	233
68	13	9	34	249	94	29	123	497	79	24	247
70	11	9	35	250	39	17	124	509	116	25	253
71	38	16	35	253	141	30	125	513	112	25	255
87	13	10	43	263	61	18	130	514	139	41	256
100	29	27	50	275	49	18	137	517	108	25	257
101	21	18	50	294	67	19	146	521	201	42	259
102	19	11	51	297	64	19	147	532	93	25	265
103	18	11	51	298	13	19	148	532	309	85	265
105	16	11	52	298	67	31	148	537	337	64	266
106	37	19	52	301	176	64	150	547	295	44	271
113	31	12	56	303	115	32	150	555	121	26	276
125	19	12	62	311	168	49	154	571	202	66	283
134	39	21	67	312	49	19	155	578	151	27	287
137	30	13	68	316	65	33	156	583	93	26	290
138	31	13	68	329	71	20	163	599	139	44	298
139	30	13	69	337	188	51	167	610	351	91	304
141	37	22	70	353	97	34	176	625	241	46	310
145	51	44	72	376	145	71	187	633	151	28	314
148	85	23	73	379	159	36	188	638	93	47	316
159	37	14	79	383	101	22	190	673	473	72	334
179	39	24	89	386	211	37	191	676	181	47	337
187	69	50	93	411	73	22	204	706	135	29	351
191	34	15	95	424	157	37	211	709	251	49	352
197	51	26	97	428	89	23	212	710	131	29	353
217	39	16	108	441	121	38	219	711	493	74	353

Table 2: Lens spaces with $712 \leq p \leq 2007$ which homology sphere with $d(Y) = 2$ yield.

p	q	h	g	p	q	h	g	p	q	h	g
715	199	98	356	1103	291	60	550	1552	849	145	774
736	393	51	365	1129	240	37	562	1563	640	73	778
739	161	30	367	1135	234	37	565	1583	334	110	787
767	133	30	382	1141	421	62	568	1618	149	75	804
773	181	50	385	1162	253	125	579	1634	427	73	815
789	172	31	392	1163	149	38	578	1641	340	112	816
790	171	31	393	1168	201	37	582	1653	283	44	824
796	165	31	396	1171	321	63	582	1717	307	152	856
805	211	104	401	1173	814	95	583	1727	389	46	860
813	211	32	404	1191	253	38	593	1742	283	45	868
823	340	53	409	1198	631	65	595	1758	451	47	875
828	133	31	412	1223	848	97	608	1772	925	79	881
841	107	54	417	1226	257	63	611	1779	337	46	886
873	151	32	435	1243	201	38	619	1783	427	76	889
878	129	33	436	1276	291	131	636	1803	406	47	898
893	237	54	445	1285	336	66	639	1807	309	46	901
919	379	56	457	1298	223	39	647	1811	1247	118	901
925	519	112	461	1331	135	68	661	1841	561	78	917
938	151	33	467	1376	361	67	686	1849	360	47	921
953	505	58	473	1377	223	40	686	1853	189	48	922
975	181	34	485	1379	302	41	686	1855	319	158	925
991	265	87	492	1403	361	42	698	1857	352	47	925
999	226	35	497	1408	273	41	701	1873	1289	120	932
1004	233	57	500	1414	267	41	704	1887	406	80	939
1021	301	58	508	1426	783	139	711	1900	289	47	947
1027	189	35	511	1437	589	70	715	1933	163	82	961
1027	573	118	512	1447	317	42	720	1963	511	80	979
1033	192	35	514	1471	771	72	731	1985	416	49	989
1037	271	89	515	1488	169	43	740	1993	408	49	993
1057	239	36	526	1513	285	43	760	2001	721	82	997
1072	121	61	532	1526	323	43	760				
1088	281	37	541	1534	315	43	764				

Conjecture 4.1 Suppose $L(p, q) = \Sigma(2, 3, 5)_p(K)$. For $h \in \mathcal{H}(p, K)$, either of the following 6 patterns holds:

1. $L(p, q) = L(54\ell^2 + 15\ell + 1, 27\ell^2 + 21\ell + 3)$ for $\ell \in \mathbb{Z} \setminus \{0\}$,
2. $L(p, q) = L(54\ell^2 + 39\ell + 7, 27\ell^2 + 33\ell + 9)$ for $\ell \in \mathbb{Z} \setminus \{0\}$,
3. $L(p, q) = L(69\ell^2 + 17\ell + 1, 46\ell^2 + 19\ell + 2)$ for $\ell \in \mathbb{Z} \setminus \{0\}$,
4. $L(p, q) = L(69\ell^2 + 29\ell + 3, 46\ell^2 + 27\ell + 4)$ for $\ell \in \mathbb{Z} \setminus \{0\}$,
5. $3.21 \leq \frac{h^2}{p} \leq 3.61$,
6. $1.15 \leq \frac{h^2}{p} \leq 1.28$.

Acknowledgments

The author would like express the gratitude for Professor Masaaki Ue and would thank for T. Kadokami's organizing seminars and useful communication there with N. Maruyama and T. Sakai as well. The author is grateful for J. Rasmussen's pointing out mistakes of the list in Lemma 4.1. The author was supported by the Grant-in-Aid for JSPS Fellows for Young Scientists(17-1604).

References

- [1] J. Berge, *Some knots with surgeries yielding lens spaces*, unpublished manuscript.
- [2] R. Fintushel and R. Stern, *Constructing lens spaces by surgery on knots*, Math. Z. Vol. 175, no. 1 February, (1980), 33-51
- [3] H. Goda and M. Teragaito, *Dehn surgeries on knots which yield lens spaces and genera of knots*, Math. Proc. Cambridge Philos. Soc. 129 (2000), 501-515.
- [4] T. Kadokami, *On the Alexander polynomial satisfying Ozsváth Szabó's condition for lens surgery*, preprint
- [5] T. Kadokami, and Y. Yamada, *A deformation of the Alexander polynomials of knots yielding lens spaces*, Bull. of Austral. Math. Soc.
- [6] P. Kronheimer, T. Mrowka, P. Ozsváth, and Z. Szabó, *Monopoles and lens space surgeries*, arXiv:math.GT/0310164
- [7] K. Ichihara, T. Saito, and M. Teragaito, *Alexander polynomials of doubly primitive knots*, Proc. Amer. Math. Soc. 135 (2007), 605-615
- [8] P. Ozsváth and Z. Szabó, *Absolutely graded Floer homologies and intersection forms for four-manifolds with boundary*, Adv. Math. 173 (2003), no. 2, 179-261.
- [9] P. Ozsváth and Z. Szabó, *Holomorphic disks and genus bounds*, Geometry & Topology 8(2004) 311-334.
- [10] P. Ozsváth and Z. Szabó, *On knot Floer homology and lens surgery*, Topology Volume 44, Issue 6, November 2005, Pages 1281-1300
- [11] G. Perelman, *The entropy formula for the Ricci flow and its geometric applications* arXiv:math/0211159
- [12] J. Rasmussen, *Lens space surgeries and a conjecture of Goda and Teragaito* Geometry & Topology Vol.8(2004) 1013-1031
- [13] R. Rustamov, *Surgery formula for the renormalized Euler characteristic of Heegaard Floer homology*, arXiv:math.GT/0409294

- [14] T. Saito, *Dehn surgery and $(1,1)$ -knots in lens spaces*, Topology Appl. 154 (2007), no. 7, 1502-1515
- [15] M. Tange, *On the non-existence of lens space surgery structure*, arXiv:0707.0197
- [16] M. Tange, *Ozsváth-Szabó's correction term of lens surgery*, preprint.
- [17] M. Tange, *The dual knot of lens surgery and Alexander polynomial*, preprint.
- [18] Y. Ni, *Knot Floer homology detects fibered knots*, arXiv:math.GT/0607156




Figure 1: $h - p$ graph in $\Sigma(2,3,5)$ case.




Figure 2: $h - p$ graph in S^3 case.